

The Equivariant Chow Ring of $\mathrm{SO}(4)$

Rahul Pandharipande¹

4 July 1996

0. Introduction

Let $\mathbf{G}$ be a reductive algebraic group. The algebraic analogue of $E\mathbf{G}$ is attained by approximation. Let V be a $\mathbb{C}$ -vector space. Let $\mathbf{G} \times V \rightarrow V$ be an algebraic representation of $\mathbf{G}$. Let $W \subset V$ be a $\mathbf{G}$ -invariant open set satisfying:

- (i) The complement of W in V is of codimension greater than q .
- (ii) $\mathbf{G}$ acts freely on W .
- (iii) There exists a geometric quotient $W \rightarrow W/\mathbf{G}$.

W is an approximation of $E\mathbf{G}$ up to codimension q . Let $e = \dim(W/\mathbf{G})$. The equivariant Chow groups of $\mathbf{G}$ (acting on a point) are defined by:

$$(1) \quad A_{-j}^{\mathbf{G}}(\text{point}) = A_{e-j}(W/\mathbf{G})$$

for $0 \leq j \leq q$. An argument is required to check the Chow groups are well-defined (see [EG1]). The basic properties of equivariant Chow groups are established in [EG1]. In particular, there is a natural intersection ring structure on $A_i^{\mathbf{G}}(\text{point})$. For notational convenience, a superscript will denote the Chow group codimension:

$$A_{-j}^{\mathbf{G}}(\text{point}) = A_{\mathbf{G}}^j(\text{point}).$$

Equation (1) becomes:

$$\forall 0 \leq j \leq q, \quad A_{\mathbf{G}}^j(\text{point}) = A^j(W/\mathbf{G}).$$

$W/\mathbf{G}$ is an approximation of $B\mathbf{G}$. $A_{\mathbf{G}}^*(\text{point})$ is called the (equivariant) Chow ring of $\mathbf{G}$. $A_{\mathbf{G}}^*(\text{point})$ is naturally isomorphic to the ring of algebraic characteristic classes of (étale locally trivial) principal $\mathbf{G}$ -bundles. The equivariant Chow ring of $\mathbf{G}$ was first defined by B. Totaro in [T].

Consider now the orthogonal and special orthogonal algebraic groups (over $\mathbb{C}$). The Chow ring of $\mathbf{O}(n)$ is generated by the Chern classes of the standard representation. The odd classes are 2-torsion:

$$A_{\mathbf{O}(n)}^*(\text{point}) = \mathbb{Z}[c_1, \dots, c_n]/(2c_1, 2c_3, 2c_5, \dots).$$

The Chow ring of $\mathbf{SO}(n = 2k + 1)$ is also generated by the Chern classes of the standard representation. The odd classes are 2-torsion and $c_1 = 0$:

$$A_{\mathbf{SO}(n=2k+1)}^*(\text{point}) = \mathbb{Z}[c_1, \dots, c_n]/(c_1, 2c_3, 2c_5, \dots).$$

¹Research partially supported by an NSF Post-Doctoral Fellowship.

$A_{\mathbf{O}(n)}^*(\text{point})$ was first computed by B. Totaro. Algebraic computations of $A_{\mathbf{O}(n)}^*(\text{point})$ and $A_{\mathbf{SO}(2k+1)}^*(\text{point})$ can be found in [P2]. The Chow ring of $\mathbf{SO}(n)$ has been computed with $\mathbb{Q}$ -coefficients in [EG2]. The integral Chow ring of $\mathbf{SO}(n = 2k)$ is not known in general.

Since $\mathbf{BO}(n)$ is approximated by the set of non-degenerate quadratic forms in $\text{Sym}^2 S^*$ (where $S \rightarrow \mathbf{G}(n, \infty)$ is the tautological sub-bundle), the Chow ring can be analyzed by degeneracy calculations ([P2]). Algebraic $\mathbf{BSO}(n)$ double covers $\mathbf{BO}(n)$. If $n = 2k + 1$, there is a product decomposition $\mathbf{O}(n) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbf{SO}(n)$. As a result, the double cover geometry for $n = 2k + 1$ is tractable and a computation of $A_{\mathbf{SO}(2k+1)}^*(\text{point})$ can be made. In case $n = 2k$, the double cover geometry is more complicated.

Since $\mathbf{SO}(2) \cong \mathbb{C}^*$, the first non-trivial even case is $\mathbf{SO}(4)$. In this paper, the ring $A_{\mathbf{SO}(4)}^*(\text{point})$ is determined. Let c_1, c_2, c_3, c_4 be Chern classes of the standard representation of $\mathbf{SO}(4)$. Let F be one of the two distinct irreducible 3-dimensional representations of $\mathbf{SO}(4)$, and let f_2 be the second Chern class F .

Theorem 1. $A_{\mathbf{SO}(4)}^*(\text{point})$ is generated by the Chern classes c_1, c_2, c_3, c_4 , and f_2 . Define $x \in A_{\mathbf{SO}(4)}^2$ by $x = c_2 - f_2$.

$$A_{\mathbf{SO}(4)}^*(\text{point}) = \mathbb{Z}[c_1, c_2, c_3, c_4, x] / (c_1, 2c_3, xc_3, x^2 - 4c_4)$$

Let $\tilde{F}$ be the other irreducible 3-dimensional representation of $\mathbf{SO}(4)$. Let $\tilde{f}_2$ be the second Chern class of $\tilde{F}$. Since (see [FH])

$$F \oplus \tilde{F} \cong \wedge^2 V,$$

the relation $\tilde{f}_2 = 2c_2 - f_2$ is obtained. Hence, $c_2 - \tilde{f}_2 = -x$. The presentation in Theorem 1 does not depend upon the choice of 3-dimensional representation.

Thanks are due to D. Edidin, W. Fulton, W. Graham, and B. Totaro for conversations about $\mathbf{BSO}(n)$. The $\mathbf{SO}(4)$ calculation presented here is similar in spirit to the $\mathbf{SO}(2k + 1)$ calculations of [P2]. In [P1] and [P2], equivariant Chow rings are used to compute ordinary Chow rings of certain moduli spaces of maps and Hilbert schemes of rational curves.

1. Ruled Quadric Surfaces

Let $V \cong \mathbb{C}^4$ be equipped with a non-degenerate quadratic form Q . A *ruled quadric surface* in $\mathbf{P}(V)$ is a pair (X, r) where $X \subset \mathbf{P}(V)$ is a nonsingular quadric surface and r is a choice of ruling. Let $\mathcal{X} \subset$

$\mathbf{P}(\mathrm{Sym}^2 V^*)$ be the parameter space of nonsingular quadrics. The parameter space of ruled quadrics, $\mathcal{X}_{\text{ruled}}$, is an étale double cover of $\mathcal{X}$ via the natural map: $\mathcal{X}_{\text{ruled}} \rightarrow \mathcal{X}$.

There are natural maps $\mathbf{SO}(V) \rightarrow \mathbf{PSO}(V) \subset \mathbf{PGL}(V)$ and $\mathbf{O}(V) \rightarrow \mathbf{PO}(V) \subset \mathbf{PGL}(V)$. Let $\mathbf{SO}(V)$ and $\mathbf{O}(V)$ act on $\mathbf{PGL}(V)$ on the right via these maps. There exist geometric quotients (see [P2]):

$$\mathbf{PGL}(V)/\mathbf{SO}(V) \rightarrow \mathbf{PGL}(V)/\mathbf{O}(V).$$

Consider the quadric surface $(Q) \subset \mathbf{P}(V)$ obtained from the quadratic form. The standard left action $\mathbf{PGL}(V) \times \mathbf{P}(V) \rightarrow \mathbf{P}(V)$ yields a transitive $\mathbf{PGL}(V)$ -action on the space of nonsingular quadric surfaces. The stabilizer of (Q) for this action is exactly $\mathbf{PO}(V) \subset \mathbf{PGL}(V)$. Hence, there is a canonical isomorphism

$$\mathbf{PGL}(V)/\mathbf{O}(V) \cong \mathcal{X}.$$

For the entire paper, fix a ruling r of (Q) . Since $\mathbf{PGL}(V)$ acts transitively on the space of ruled quadrics and the stabilizer of $((Q), r)$ is exactly $\mathbf{PSO}(V) \subset \mathbf{PGL}(V)$, there is a canonical isomorphism determined by $((Q), r)$:

$$\mathbf{PGL}(V)/\mathbf{SO}(V) \cong \mathcal{X}_{\text{ruled}}.$$

There is a canonical Plücker embedding $\mathbf{G}(2, V) \hookrightarrow \mathbf{P}(\wedge^2 V)$. Let $Z \subset \mathbf{G}(3, \wedge^2 V)$ be the open locus of 2-planes in $\mathbf{P}(\wedge^2 V)$ which intersect $\mathbf{G}(2, V)$ transversely in a nonsingular conic curve.

Lemma 1. *There is a canonical isomorphism $Z \cong \mathcal{X}_{\text{ruled}}$.*

Proof. The family of lines determined by a nonsingular plane conic $C \subset \mathbf{G}(2, V) \subset \mathbf{P}(\wedge^2 V)$ sweeps out an irreducible degree 2 surface in $\mathbf{P}(V)$. There are three possibilities for this degree 2 surface: a double plane, a quadric cone, or a nonsingular quadric surface. If the conic C sweeps out a double plane $H \subset \mathbf{P}(V)$, then $C \subset P \subset \mathbf{G}(2, V)$ where P is the plane of all lines contained in H . If C sweeps out a quadric cone, then $C \subset P \subset \mathbf{G}(2, V)$ where P is the plane of all lines passing through the vertex of the cone. Hence, if C is the transverse intersection $P \cap \mathbf{G}(2, V)$ of a plane, then C must correspond to a ruling of a unique nonsingular quadric surface. Conversely, a ruling of a nonsingular quadric surface yields a conic curve in $\mathbf{G}(2, V)$ which is the transverse intersection of a unique 2-plane in $\mathbf{P}(V)$. These maps are easily seen to be algebraic. $\square$

By Lemma 1, the ruled quadric $((Q), r)$ corresponds to a 3-dimensional subspace $F \subset \wedge^2 V$. F is $\mathbf{SO}(V)$ -invariant since $((Q), r)$ is stabilized by $\mathbf{SO}(V)$. F is therefore a 3-dimensional representation of $\mathbf{SO}(V)$. Let

s be the *other* ruling of (Q) . $((Q), s)$ similarly corresponds to an invariant 3-dimensional subspace $\tilde{F} \subset \wedge^2 V$. The $\mathbf{SO}(V)$ representation $\wedge^2 V$ decomposes as $\wedge^2 V \cong F \oplus \tilde{F}$.

2. $BSO(V)$

Let $V \cong \mathbb{C}^4$ be equipped with a non-degenerate quadratic form as before. Approximations to $ESO(V)$ and $BSO(V)$ are obtained via direct sums of the representation V^* . Let $m \gg 0$ and let

$$W_m \subset \oplus_1^m V^*$$

denote the spanning locus. W_m is the locus of m -tuples of vectors of V^* which span V^* . The natural action of $\mathbf{SO}(V)$ on W_m is free and has a geometric quotient (see [P2]). The codimension of the complement of W_m in $\oplus_1^m V^*$ is $m - 3$. W_m is an approximation of $ESO(V)$ up to codimension $m - 4$. Therefore

$$BSO(V) = m \xrightarrow{Lim} \infty W_m / \mathbf{SO}(V),$$

$$A_{\mathbf{SO}(V)}^*(\text{point}) = m \xrightarrow{Lim} \infty A^*(W_m / \mathbf{SO}(V)).$$

There is a scalar $\mathbb{C}^*$ -action on W_m . Let $\mathbf{P}(W_m) = W_m / \mathbb{C}^*$. Since this $\mathbb{C}^*$ -action commutes with the $\mathbf{SO}(V)$ -action, there is diagram of quotients:

$$(2) \quad \begin{array}{ccc} W_m & \xrightarrow{\tau_1} & W_m / \mathbf{SO}(V) \\ i_1 \downarrow & & \downarrow i_2 \\ \mathbf{P}(W_m) = W_m / \mathbb{C}^* & \xrightarrow{\tau_2} & \mathbf{P}(W_m) / \mathbf{SO}(V) \end{array}$$

All the maps in (2) are quotient maps:

- (i) i_1 is a free $\mathbb{C}^*$ -quotient.
- (ii) i_2 is a free $\mathbb{C}^*/(\pm)$ -quotient.
- (iii) τ_1 is a free $\mathbf{SO}(V)$ -quotient.
- (iv) τ_2 is a free $\mathbf{PSO}(V)$ -quotient.

The existence of these quotients is easily deduced (see [P2]).

First consider the space $\mathbf{P}(W_m) / \mathbf{SO}(V)$. Let Q be the quadratic form on V and let r be the ruling of the quadric surface $(Q) \subset \mathbf{P}(V)$ fixed in section 1. An element $f \in \mathbf{P}(W_m)$ yields a canonical embedding

$$\mu_f : \mathbf{P}(V) \hookrightarrow \mathbf{P}(\mathbb{C}^m).$$

The image under μ_f of $((Q), r)$ is a ruled quadric surface in $\mathbf{P}(\mathbb{C}^m)$ associated canonically to $f \in \mathbf{P}(W_m)$. Since $\mathbf{PSO}(V) \subset \mathbf{PGL}(V)$ is exactly the stabilizer of the ruled quadric $((Q), r)$, it follows that $\mathbf{P}(W_m) / \mathbf{SO}(V)$ is isomorphic to the parameter space of ruled quadric

surfaces in $\mathbf{P}(\mathbb{C}^m)$. Since a ruled quadric surface in $\mathbf{P}(\mathbb{C}^m)$ spans a unique 3-plane in $\mathbf{P}(\mathbb{C}^m)$, the parameter space is fibered over $\mathbf{G}(4, m)$. By Lemma 1, the parameter space of ruled quadric surface in $\mathbf{P}(\mathbb{C}^m)$ is canonically isomorphic to an open set

$$Z \subset \mathbf{G}(3, \wedge^2 S)$$

where $S \rightarrow \mathbf{G}(4, m)$ is the tautological sub-bundle.

The Chow computations in section 3 will require two results about line bundles. We have seen $\mathbf{P}(W_m)/\mathbf{SO}(V)$ is canonically fibered over $\mathbf{G}(4, m)$. Let c_1 be the first Chern class of the tautological bundle S on $\mathbf{G}(4, m)$. Let c_1 also denote the pull-back of this class to $\mathbf{P}(W_m)/\mathbf{SO}(V)$. For $m > 4$, $A^1(\mathbf{P}(W_m)) \cong \mathbb{Z}$ with generator $c_1(\mathcal{O}_{\mathbf{P}}(-1))$ (which is the Chern class of the line bundle associated to the $\mathbb{C}^*$ -bundle i_1 .)

Lemma 2. $\tau_2^*(c_1) = c_1(\mathcal{O}_{\mathbf{P}}(-4))$.

Proof. Elements of $\mathbf{P}(W_m)$ correspond to embeddings of $\mathbf{P}(V)$ in $\mathbf{P}(\mathbb{C}^m)$. The class $\tau_2^*(-c_1)$ is the divisor class of embeddings that meet a fixed $(m-5)$ -plane in $\mathbf{P}(\mathbb{C}^m) = \mathbf{P}^{m-1}$. This divisor class is determined by a 4×4 determinant. Hence, $\tau_2^*(-c_1) = c_1(\mathcal{O}_{\mathbf{P}}(4))$. $\square$

The map $i_2 : W_m/\mathbf{SO}(V) \rightarrow \mathbf{P}(W_m)/\mathbf{SO}(V)$ is a $\mathbb{C}^*/(\pm)$ -bundle. Since there is an abstract isomorphism $\mathbb{C}^*/(\pm) \cong \mathbb{C}^*$, i_2 is also a $\mathbb{C}^*$ -bundle. Let N be the line bundle on $\mathbf{P}(W_m)/\mathbf{SO}(V)$ canonically associated to i_2 .

Lemma 3. $\tau_2^*(N) \cong \mathcal{O}_{\mathbf{P}}(-2)$.

Proof. Let $i_1/(\pm) : W_m/(\pm) \rightarrow \mathbf{P}(W_m)$. The map $i_1/(\pm)$ is a free $\mathbb{C}^*/(\pm)$ -quotient. The line bundle associated to the $\mathbb{C}^*/(\pm)$ -bundle $i_1/(\pm)$ is $\mathcal{O}_{\mathbf{P}}(-2)$. The map $\tau_1/(\pm) : W_m/(\pm) \rightarrow W_m/\mathbf{SO}(V)$ is $\mathbb{C}^*/(\pm)$ -equivariant. Hence, $\tau_2^*(N) \cong \mathcal{O}_{\mathbf{P}}(-2)$. $\square$

3. Chow Calculations

In this section, the Chow ring of $W_m/\mathbf{SO}(V)$ is determined (up to codimension $m-4$). Consider the parameter space of ruled quadrics in $\mathbf{P}(\mathbb{C}^m)$:

$$Z \subset \mathbf{G}(3, \wedge^2 S) \rightarrow \mathbf{G}(4, m).$$

Let D be the complement of Z in $\mathbf{G}(3, \wedge^2 S)$. Following the notation of section 2, $W_m/\mathbf{SO}(V)$ is the $\mathbb{C}^*$ -bundle associated to a line bundle $N \rightarrow Z$. Therefore,

$$A^*(W_m/\mathbf{SO}(V)) \cong A^*(Z)/(c_1(N)) \cong A^*(\mathbf{G}(3, \wedge^2 S))/(I_D, c_1(\overline{N}))$$

where $I_D \subset A^*(\mathbf{G}(3, \wedge^2 S))$ is the ideal generated by cycles supported on D and $\overline{N}$ is any extension of N to $\mathbf{G}(3, \wedge^2 S)$. The ideal I_D is determined by constructing a well-behaved variety which surjects onto D .

Let $\mathbf{G}(2, S) \hookrightarrow \mathbf{P}(\wedge^2 S)$ be the canonical relative Plücker embedding. D is exactly the locus of 2-planes in the fibers of $\mathbf{P}(\wedge^2 S)$ which do not meet $\mathbf{G}(2, S)$ transversely in a nonsingular conic curve. Equivalently, D is the locus of 2-planes P in the fibers of $\mathbf{P}(\wedge^2 S)$ which satisfy one of the following conditions:

- (i) $P \cap \mathbf{G}(2, S)$ is a pair of distinct lines in P .
- (ii) $P \cap \mathbf{G}(2, S)$ is a double line in P .
- (iii) $P \cap \mathbf{G}(2, S) = P$.

D is dominated by a canonical Grassmannian bundle over $\mathbf{G}(2, S)$. Let $B \rightarrow \mathbf{G}(2, S)$ be the tautological sub-bundle. By wedging, there is canonical surjective bundle map on $\mathbf{G}(2, S)$:

$$\wedge^2 S \otimes \wedge^2 B \rightarrow \wedge^4 S$$

which induces a canonical sequence on $\mathbf{G}(2, S)$:

$$(3) \quad 0 \rightarrow K \rightarrow \wedge^2 S \rightarrow \wedge^4 S \otimes (\wedge^2 B)^* \rightarrow 0.$$

There is a canonical inclusion $\wedge^2 B \subset K$ and a quotient sequence

$$(4) \quad 0 \rightarrow \wedge^2 B \rightarrow K \rightarrow E \rightarrow 0$$

on $\mathbf{G}(2, S)$. The geometric interpretation of these sequences is as follows. Let $\xi \in \mathbf{G}(2, S)$. $\mathbf{P}(K_\xi) \subset \mathbf{P}(\wedge^2 S_\xi)$ is the projective tangent space to $\mathbf{G}(2, S_\xi)$ at ξ . $\mathbf{P}(\wedge^2 B_\xi)$ in $\mathbf{P}(\wedge^2 S_\xi)$ is the Plücker image of the point ξ . The fiber of the Grassmannian bundle

$$\mathbf{G}(2, E) \rightarrow \mathbf{G}(2, S)$$

over ξ corresponds to the 2-planes P of $\mathbf{P}(\wedge^2 S_\xi)$ that are tangent to $\mathbf{G}(2, S)$ at ξ . There is a canonical map

$$\rho : \mathbf{G}(2, E) \rightarrow D$$

which is a surjection of algebraic varieties. Let $[P] \in D$. The fiber of ρ over $[P]$ is simply the set of points of $P \cap \mathbf{G}(2, S)$ where P is tangent to $\mathbf{G}(2, S)$. In case (i) above, the fiber is a point. In case (ii), the fiber is a straight line in $\mathbf{P}(\wedge^2 S)$. In case (iii), the fiber is 2-plane in $\mathbf{P}(\wedge^2 S)$. Hence, there is stratification of D by intersection type (i-iii) where ρ is a projective bundle over each stratum. The Chow groups of $\mathbf{G}(2, E)$ therefore surject upon the Chow groups of D .

The ideal I_D is determined by calculating the push-forwards of the Chow classes of $\mathbf{G}(2, E)$ to $\mathbf{G}(3, \wedge^2 S)$. Consider the projection

$$\pi : \mathcal{G} = \mathbf{G}(3, \wedge^2 S) \times_{\mathbf{G}(4, m)} \mathbf{G}(2, S) \rightarrow \mathbf{G}(3, \wedge^2 S).$$

The sequences (3) and (4) pull-back to $\mathcal{G}$. Let $F \rightarrow \mathbf{G}(3, \wedge^2 S)$ denote the tautological sub-bundle (and also let F denote the pull-back to $\mathcal{G}$ of this bundle). There is a canonical inclusion

$$\iota : \mathbf{G}(2, E) \hookrightarrow \mathcal{G}$$

determined by the sequences (3) and (4). $\mathbf{G}(2, E) \subset \mathcal{G}$ is the closed subvariety of points $g \in \mathcal{G}$ where

$$\wedge^2 B_g \subset F_g \subset K_g.$$

The class $[\mathbf{G}(2, E)]$ in Chow ring of $A^*(\mathcal{G})$ is easily found by degeneracy calculations. Let c_1, c_2, c_3, c_4 be the Chern classes of $S \rightarrow \mathbf{G}(4, m)$. Let b_1, b_2 be the Chern classes of $B \rightarrow \mathbf{G}(2, S)$. Let f_1, f_2, f_3 be the Chern classes of $F \rightarrow \mathbf{G}(3, \wedge^2 S)$. Since $\mathcal{G}$ is a tower of Grassmannian bundles, these Chern classes c_i, b_j, f_k generate $A^*(\mathcal{G})$. Let Y be the locus of points $g \in \mathcal{G}$ such that $F_g \subset K_g$. Y is the nonsingular degeneracy locus of the canonical bundle map on $\mathcal{G}$,

$$F \rightarrow \wedge^4 S \otimes (\wedge^2 B)^*,$$

obtained from the inclusion $F \subset \wedge^2 S$ and sequence (3). By the Thom-Porteous formula on $\mathcal{G}$ (see [F]),

$$\begin{aligned} A^*(\mathcal{G}) \ni [Y] &= c_3(F^* \otimes \wedge^4 S \otimes (\wedge^2 B)^*) \\ &= -f_3 + (c_1 - b_1)f_2 - (c_1 - b_1)^2 f_1 + (c_1 - b_1)^3. \end{aligned}$$

Y is canonically isomorphic to the Grassmannian bundle $\mathbf{G}(3, K) \rightarrow \mathbf{G}(2, S)$. There is natural bundle quotient sequence on Y :

$$(5) \quad 0 \rightarrow F \rightarrow K \rightarrow K/F \rightarrow 0.$$

The locus $\mathbf{G}(2, E) \subset Y$ is the set of points $y \in Y$ such that $\wedge^2 B_y \subset F_y$. $\mathbf{G}(2, E) \subset Y$ is the nonsingular degeneracy locus of the canonical bundle map on Y ,

$$\wedge^2 B \rightarrow K/F,$$

obtained from the sequences (4) and (5). By the Thom-Porteous formula on Y ,

$$\begin{aligned} A^*(Y) \ni [\mathbf{G}(2, E)] &= c_2((K/F) \otimes (\wedge^2 B)^*) \\ &= b_1^2 - c_1 b_1 + c_1^2 - 2c_1 f_1 + f_1^2 - f_2 + 2c_2. \end{aligned}$$

The class $[\mathbf{G}(2, E)] \in A^*(\mathcal{G})$ is there expressed by

$$\begin{aligned} A^*(\mathcal{G}) \ni [\mathbf{G}(2, E)] &= (-f_3 + (c_1 - b_1)f_2 - (c_1 - b_1)^2 f_1 + (c_1 - b_1)^3) \\ &\quad \cdot (b_1^2 - c_1 b_1 + c_1^2 - 2c_1 f_1 + f_1^2 - f_2 + 2c_2). \end{aligned}$$

Since $\mathbf{G}(2, E)$ is a Grassmannian bundle over $\mathbf{G}(2, S)$, the Chow ring of $\mathbf{G}(2, E)$ is generated over $A^*(\mathbf{G}(2, S))$ by the Chern classes h_1, h_2 of the tautological sub-bundle $H \rightarrow \mathbf{G}(2, E)$. Via the embedding $\iota : \mathbf{G}(2, E) \hookrightarrow \mathcal{G}$, H is isomorphic to $\iota^*(F)/\iota^*(\wedge^2 B)$. The Chern classes h_1 and h_2 can be expressed via ι in terms of the classes b_j and f_k . Therefore, the classes

$$\mathbf{G}(2, E) \cap M(c_1, c_2, c_3, c_4, b_1, b_2, f_1, f_2, f_3)$$

(where M is monomial in the Chern classes) span the Chow ring of $\mathbf{G}(2, E)$. The ideal $I_D \subset A^*(\mathbf{G}(3, \wedge^2 S))$ is generated by the π push-forwards of the classes (6):

$$(6) \quad [\mathbf{G}(2, E)] \cdot M(c_1, c_2, c_3, c_4, b_1, b_2, f_1, f_2, f_3) \in A^*(\mathcal{G})$$

Since the classes c_i, f_k in $A^*(\mathcal{G})$ are pull-backs from $\mathbf{G}(3, \wedge^2 S)$, I_D is generated by the elements

$$\pi_*([\mathbf{G}(2, E)] \cdot M(b_1, b_2)).$$

By the standard relations satisfied by the classes b_1 and b_2 over $A^*(\mathbf{G}(4, m))$, it follows that I_D is generated by:

$$\begin{aligned} &\pi_*([\mathbf{G}(2, E)]) \\ &\pi_*([\mathbf{G}(2, E)] \cdot b_1) \\ &\pi_*([\mathbf{G}(2, E)] \cdot b_1^2) \quad \pi_*([\mathbf{G}(2, E)] \cdot b_2) \\ &\pi_*([\mathbf{G}(2, E)] \cdot b_1 b_2) \\ &\pi_*([\mathbf{G}(2, E)] \cdot b_1^2 b_2). \end{aligned}$$

Since the class $[\mathbf{G}(2, E)]$ is determined explicitly in $A^*(\mathcal{G})$, these six push-forwards can be easily computed by hand or by the the Maple package Schubert ([KS]). The first push-forward is:

$$\pi_*([\mathbf{G}(2, E)]) = 13c_1 - 2f_1.$$

Lemma 4. *The pair $(13c_1 - 2f_1, c_1(\overline{N}))$ generates $A^1(\mathbf{G}(3, \wedge^2 S))$.*

Proof. Recall the notation of diagram (2):

$$\tau_2 : \mathbf{P}(W_m) \rightarrow Z \subset \mathbf{G}(3, \wedge^2 S).$$

Let $L = c_1(\mathcal{O}_{\mathbf{P}}(-1))$ be a generator of $A^1(\mathbf{P}(W_m))$. By Lemma 2, $\tau_2^*(c_1) \stackrel{\sim}{=} 4L$. Since $\tau_2^*([D]) = \tau_2^*(13c_1 - 2f_1) = 0$, $\tau_2^*(f_1) = 26L$. Therefore the image of τ_2^* is the subgroup $\mathbb{Z}(2L)$.

Since $[D] = 13c_1 - 2f_1$ is not divisible in $A^1(\mathbf{G}(3, \wedge^2 S))$, $A^1(Z) \cong \mathbb{Z}$ and τ_2^* is an isomorphism:

$$\tau_2^* : A^1(Z) \xrightarrow{\sim} \mathbb{Z}(2L).$$

By Lemma 3, $\tau_2^*(c_1(N)) = 2L$. Therefore, $c_1(N)$ generates $A^1(Z)$. It now follows that the pair $(13c_1 - 2f_1, c_1(\overline{N}))$ generates the group $A^1(\mathbf{G}(3, \wedge^2 S))$. $\square$

Therefore, $(I_D, c_1(\overline{N})) = (I_D, c_1, f_1)$.

By Lemma 4, it suffices to compute the five remaining push-forwards modulo the ideal $J = (c_1, f_1)$. The results are (modulo J):

$$\begin{aligned} \pi_*([\mathbf{G}(2, E)] \cdot b_1) &= 0. \\ \pi_*([\mathbf{G}(2, E)] \cdot b_1^2) &= -2f_3. \\ \pi_*([\mathbf{G}(2, E)] \cdot b_2) &= c_3 - f_3. \\ \pi_*([\mathbf{G}(2, E)] \cdot b_1 b_2) &= (c_2 - f_2)^2 - 4c_4. \\ \pi_*([\mathbf{G}(2, E)] \cdot b_1^2 b_2) &= c_2 f_3 + f_2 c_3. \end{aligned}$$

Hence $(I_D, c_1(\overline{N})) = (c_1, f_1, 2c_3, c_3 - f_3, (c_2 - f_2)^2 - 4c_4, (c_2 - f_2)c_3)$.

The ring $A^*(\mathbf{G}(4, m))$ is freely generated (up to codimension $m - 4$) by c_1, c_2, c_3, c_4 . The ring $A^*(\mathbf{G}(3, \wedge^2 S))$ has the following presentation (up to codimension $m - 4$):

$$A^*(\mathbf{G}(3, \wedge^2 S)) \cong \mathbb{Z}[c_1, c_2, c_3, c_4, f_1, f_2, f_3]/(t_4, t_5, t_6)$$

where the t_4, t_5, t_6 are the Chern classes of the tautological quotient bundle T : $0 \rightarrow F \rightarrow \wedge^2 S \rightarrow T \rightarrow 0$. We find (modulo J):

$$\begin{aligned} t_4 &= (c_2 - f_2)^2 - 4c_4. \\ t_5 &= 2f_2 f_3 - 2c_2 f_3. \\ t_6 &= f_2(-(c_2 - f_2)^2 + 4c_4) + f_3^2 - c_3^2. \end{aligned}$$

There is a presentation: $A^*(\mathbf{G}(3, \wedge^2 S))/(I_D, c_1(\overline{N})) \cong$

$$(7) \quad \mathbb{Z}[c_1, c_2, c_3, c_4, f_1, f_2, f_3]/(I_D, c_1, f_1, t_4, t_5, t_6)$$

(up to codimension $m - 4$). Surprisingly, the relations t_4, t_5, t_6 are contained in the ideal (I_D, c_1, f_1) . By the limit procedure,

$$A_{\mathbf{SO}(4)}^*(\text{point}) \cong \mathbb{Z}[c_1, c_2, c_3, c_4, f_2]/(c_1, 2c_3, (c_2 - f_2)c_3, (c_2 - f_2)^2 - 4c_4).$$

The vector bundles $S, F \subset \wedge^2 S$ on the approximation $W_m/\mathbf{SO}(V)$ are easily seen to be obtained from the principal $\mathbf{SO}(V)$ -bundle

$$W_m \rightarrow W_m/\mathbf{SO}(V)$$

and the representations $V, F \subset \wedge^2 V$ defined in section 1. Define

$$x = c_2 - f_2.$$

Theorem 1 is proved.

REFERENCES

- [EG1] D. Edidin and W. Graham, *Equivariant intersection theory*, preprint 1996.
- [EG2] D. Edidin and W. Graham, *Characteristic classes of principal bundles in algebraic geometry*, preprint 1995.
- [F] W. Fulton, *Intersection theory*, Springer-Verlag: Berlin, 1984.
- [FH] W. Fulton and J. Harris, *Representation Theory*, Springer-Verlag: New York, 1991.
- [KS] S. Katz and S. A. Stromme, *Schubert: A Maple package for intersection theory in algebraic geometry*.
- [P1] R. Pandharipande, *The Chow ring of the non-linear Grassmannian*, preprint 1996.
- [P2] R. Pandharipande, *The Chow ring of the Hilbert scheme of rational normal curves*, preprint 1996.
- [T] B. Totaro, *The Chow ring of the symmetric group*, preprint 1994.

Department of Mathematics
University of Chicago
5734 S. University Ave. 60637
rahul@math.uchicago.edu